
Mathematical treatise: Fixed time-step issue

Angular collision response have a dynamic contact time depending on distance to the torque axis.



 ΔH : Change in angular momentum.

 \vec{r} : Offset to point of contact.

 $J_{\vec{r}}$: Particle momentum.

 $\vec{w} \& \vec{w}_{\vec{r}}$: Angular velocity.

 Δs : Collision contact stretch, relative to $M_{\Delta H}$.

 Δt_f : Fixed time periodicy of simulation.

 $\Delta t_d(\vec{r})$: Real contact duration time.

 $c_{\Delta t} = \Delta t_d / \Delta t_f$: Regulating time coefficient.

Objective

Present a formula for the $c_{\Delta t}$ coefficient, that can be implemented into a 3D physics engine with fixed time periodicy.

Calculus

$$\begin{split} c_{\mathit{At}} &= \frac{\varDelta t_{\mathit{d}}(\vec{r})}{\varDelta t_{\mathit{f}}} \;, \quad \varDelta t_{\mathit{f}} \; \text{is a constant.} \\ |\vec{w}_{\vec{r}}| &= \left\| H_{\mathsf{x}} \frac{(r_{\mathsf{y}}')^2 + (r_{\mathsf{z}}')^2}{I_{\mathit{Sxx}}} \;, \quad H_{\mathsf{y}} \frac{(r_{\mathsf{x}}')^2 + (r_{\mathsf{z}}')^2}{I_{\mathit{Syy}}} \;, \quad H_{\mathsf{z}} \frac{(r_{\mathsf{x}}')^2 + (r_{\mathsf{y}}')^2}{I_{\mathit{Szz}}} \right\|_{1} \\ |\vec{w}| &= \left\| H_{\mathsf{x}}' \frac{6}{I_{\mathit{Sxx}}} \;, \quad H_{\mathsf{y}}' \frac{6}{I_{\mathit{Syy}}} \;, \quad H_{\mathsf{z}}' \frac{6}{I_{\mathit{Szz}}} \right\|_{1} \\ \left\| \Delta s = \varDelta t_{\mathit{d}}(\vec{r}) * |\vec{\varDelta w}_{\vec{r}}| \right\|_{2} \to \left\| \varDelta t_{\mathit{d}}(\vec{r}) * |\vec{\varDelta w}_{\vec{r}}| = \varDelta t_{\mathit{f}} * |\vec{\varDelta w}| \to \frac{\varDelta t_{\mathit{d}}(\vec{r})}{\varDelta t_{\mathit{f}}} = \frac{|\vec{\varDelta w}|}{|\vec{\varDelta w}_{\vec{r}}|} = c_{\mathit{At}} \\ c_{\mathit{At}} &= \frac{|\vec{\varDelta w}|}{|\vec{\varDelta w}_{\vec{r}}|} \quad \vec{r} \;, \; \vec{\varDelta w} \; \text{and} \; \vec{\varDelta H} \; \text{ is known and} \; \vec{\varDelta w}_{\vec{r}} \; \text{ can be derived from those} \\ &= \frac{|\vec{\varDelta w}|}{|\vec{\varDelta w}_{\vec{r}}|} \quad \vec{r} \;, \; \vec{\varDelta w} \; \text{and} \; \vec{\varDelta H} \; \text{ is known and} \; \vec{\varDelta w}_{\vec{r}} \; \text{ can be derived from those} \end{split}$$

Pseudo Code

if (|vecR| > 1) vecDeltaH * = AngularImpactTimeCoefficient(|vecR, vecDeltaW, vecDeltaH);

Optimization

 $c_{At} = |\vec{\Delta w}|/|\vec{\Delta w_r}|$ can be a little bit encumbersome for a game engine. An approximative possibly good-enough alternative could be $c_{At} \sim 1/|\vec{r}|^2$.

¹ Converting angular momentum [kg*m^2/s] to angular velocity [radians/s] by Dan Andersson